

[J Agric Food Chem](#). 2013 Apr 16. [Epub ahead of print]

## **Effective Inhibition of Skin Cancer, Tyrosinase, and Antioxidative Properties by Astaxanthin and Astaxanthin Esters from the Green Alga *Haematococcus pluvialis*.**

[Rao AR](#), [Sindhuja HN](#), [Dharmesh SM](#), [Sankar KU](#), [Sarada R](#), [Ravishankar GA](#).

### **Source**

Plant Cell Biotechnology Department, ‡Biochemistry & Nutrition Department, and #Food Engineering Department, Central Food Technological Research Institute, CSIR , Mysore 570 020, Karnataka, India.

### **Abstract**

Astaxanthin mono- (AXME) and diesters (AXDE) were characterized and examined for anticancer potency with total carotenoids (TC) and astaxanthin (AX) against UV-7,12-dimethylbenz(a)anthracene (DMBA)-induced skin cancer model in rat. At 200 µg/kg bw, AXDE and AXME reduced UV-DMBA-induced tumor incidences up to 96 and 88%, respectively, when compared to AX (66%) and TC (85%). UV-DMBA has been known to generate high levels of free radicals and tyrosinase enzyme, leading to characteristic symptoms of skin pigmentation and tumor initiation. Intriguingly, ~7-fold increase in tyrosinase and 10-fold decrease in antioxidant levels were normalized by AXDE and AXME as opposed to only ~1.4-2.2-fold by AX and TC, respectively. This result together with the appearance of 72 and 58 ng/mL of retinol in the serum of respective AXE-treated (AXDE + AXME) and AX-treated animals suggested that better anticancer potency of AXEs could be due to increased bioavailability.

PMID:

23473626

[PubMed - as supplied by publisher]